

FULL PAPER

Theoretical and Experimental Methods for Studying Volumetric Behavior of Binary mixtures Containing Normal Alkanols

Mohammad Almasi


Department of Applied Chemistry, Faculty of Science, Malayer University, Malayer, 65174, Iran.

Received: 17 May 2018, Revised: 05 June 2018 and Accepted: 05 June 2018.

ABSTRACT: Molecular interactions in binary mixtures composed of a xylene and selected 1-alkanol (1-butanol up to 1-octanol) were investigated by measurement of the density at $T = 323.15$ K. From the experimental data, excess molar volumes were calculated. Obtained data were interpreted based on the type and magnitude of the physico-chemical interactions in the binary liquid systems. In this sense, PC-SAFT was used to correlate the volumetric behavior of binary mixtures. The correlated values of the model were satisfactory and the obtained data are within the uncertainty region.

KEYWORDS: Density; 1-Alkanol; PC-SAFT model.

GRAPHICAL ABSTRACT:



Excess molar volumes vs. mole fraction of xylene for binary mixtures of xylene with 1-alkanols at $T = 323.15$ K.

1. Introduction

The non-ideal behaviour of binary mixtures containing self-associated components is primarily due to the specific interactions arising from the hydrogen bond between the polar groups of the component molecules. When two liquids are mixed together, the resulting changes in the physical and thermodynamic properties can be considered as sum of several contributions

due to free volume change, change in energy, change in molecular orientations and steric hindrances [1]. Alcohols are of important class of hydrogen-bonded solvents and also of their $-\text{OH}$ group participates in hydrogen bond formation used as a donor and acceptor of electrons. Our initial studies on the thermodynamics of alcohols solutions [2-4] were motivated

by a desire to better understand the interactions among the various functional groups with alcohols. Moreover, this work contributes to test the compatibility of PC-SAFT model and also to correlate the experimental densities.

Experimental

Xylene and all 1-alkanols were analytically graded and purchased from Merck with stated level of purity greater than 99 % of mass fraction and used as received without further purification. The measured densities for pure materials were compared to the literature values. The results show that the measurements are in agreement with literature and within the experimental uncertainties. Density measurements have been performed with an Anton-Paar SVM 3000 viscometer that automatically corrects automatically viscosity related errors in the density by measuring the damping of the U-tube oscillation. Prior to each series of measurements, the apparatus was calibrated using doubly distilled degassed water and dry air at the atmospheric pressure. Pure fluids used in the binary mixtures were degasified before preparation of the samples using an ultrasonic bath. The solutions were prepared by mass, using a digital balance (Mettler AE 163, Switzerland) with the precision ± 0.01 mg and stored in airtight stopped bottles to prevent evaporation of the

samples. To minimize evaporation during the sample preparation, the less volatile component was primarily charged first. The uncertainty is 1×10^{-4} g·cm⁻³ for density measurements. Moreover, The uncertainty in the mole fraction is $\pm 1 \times 10^{-4}$.

Results and Discussion

Experimental values of densities for binary mixtures are listed in Table 1. Binary densities increase with increasing alcohols chain length. The volumetric properties are closely related to the volume change of substance, such as density. The excess molar volume can be calculated according to the following equation

$$V_m^E = V_m - \sum_{i=1}^2 x_i V_{m,i} \quad (1)$$

Where $V_{m,i}$ is the molar volume of component i and V is the volume of the mixture. The values of V_m^E can be fitted by Redlich-Kister [5] polynomial

$$V_m^E = x_1(1-x_1) \sum_{k=0}^N A_k (1-2x_1)^k \quad (2)$$

where x is the mole fraction of one component of mixture, A_k is the adjustable parameter and k is the order of polynomial. Values of excess molar volume are reported in Table 1. The measured V_m^E can be described in terms of physical, chemical and structural effects. The physical effects

consist of non-specific physical (weak) interactions, rupture of the hydrogen bonds in the self-associated molecules, weaker dipole-dipole interactions and interstitial accommodations due to different molar volumes and free volumes between unlike molecules resulting in positive V_m^E values. The chemical and structural effects which arise in accordance to the specific interactions such as formation of hydrogen bonds, charge transfer complexes, strong dipole-dipole interactions, differences in size and shape of the component molecules and make negative V_m^E values. V_m^E for xylene + 1-alkanol is positive and systematically tends to the larger positive

values with an increase in the size of 1-alkanol. This behaviour at $T= 323.15$ K is presented graphically in figure 1. The algebraic values of V_m^E fall in the order: 1-octanol > 1-heptanol > 1-hexanol > 1-pentanol > 1-butanol. When the chain length of the alcohol is increased, the hydrophobic character of the 1-alkanol is also amplified and the interactions between unlike molecules weaken, resulting in the positive contribution to V_m^E . In alcohols with higher chain length, the relative volume contributions of the polar OH versus the non polar alkyl chain for the alcohols decrease and the excess molar volume increases.

Table 1. Densities and excess molar volumes for the binary mixtures at $T= 323.15$ K and $P = 0.1$ MPa.

$T/K = 323.15$					
Densities (ρ /g·cm ⁻³) for Xylene +					
x_1	1-Butanol	1-Pentanol	1-Hexanol	1-Heptanol	1-Octanol
0	0.7860	0.7920	0.7968	0.8022	0.8037
0.081	0.7912	0.7959	0.7999	0.8045	0.8057
0.1595	0.7959	0.7996	0.8028	0.8067	0.8078
0.2395	0.8005	0.8032	0.8056	0.8086	0.8100
0.3501	0.8062	0.8077	0.8092	0.8117	0.8132
0.4397	0.8103	0.8112	0.8126	0.8142	0.8156
0.5601	0.8150	0.8158	0.8166	0.8180	0.8190
0.6497	0.8189	0.8196	0.8203	0.8211	0.8219
0.7404	0.8228	0.8244	0.8237	0.8242	0.8247
0.8492	0.8278	0.8278	0.8280	0.8282	0.8282
0.9407	0.8321	0.8319	0.8319	0.8319	0.8320
1	0.8346	0.8346	0.8346	0.8346	0.8346
V_m^E (cm ³ /mol ⁻¹)					
	0	0	0	0	0
	-0.005	-0.006	-0.003	-0.002	-0.006

-0.005	-0.006	-0.007	-0.009	-0.010
0.010	-0.002	0.004	-0.010	-0.006
0.052	0.031	0.042	0.020	0.026
0.131	0.095	0.115	0.073	0.084
0.233	0.184	0.206	0.148	0.160
0.266	0.211	0.234	0.180	0.192
0.250	0.181	0.216	0.166	0.184
0.164	0.135	0.150	0.116	0.125
0.066	0.051	0.058	0.040	0.049
0	0	0	0	0

Table 2: Binary Interaction Parameter k_{ij} and standard deviation σ for PC-SAFT Equation.

Binary Systems	k_{ij}	% σ
Xylene + 1-Butanol	0.01	1.23
Xylene + 1-Pentanol	0.014	1.02
Xylene + 1-Hexanol	0.02	1.16
Xylene + 1-Heptanol	0.012	1.14
Xylene + 1-Octanol	0.017	1.26

Our next objective in this study is to correlate binary mixtures density applying the PC-SAFT model [6, 7]. In this model, the general definition of the Helmholtz energy is illustrated by:

$$a^{res} = a^{chain} + a^{disp} + a^{assoc} \quad (3)$$

a^{chain} is the chain formation contribution computed by

$$a^{chain} = -\sum_i x_i (m_i - 1) \ln g_{ii}^{hs} \quad (4)$$

where m is the number of segments and g_{ii}^{hs} the radial distribution function for hard sphere. a^{dis} is determined by

$$a^{dis} = -2I_1 \pi m \rho^2 \left(\frac{\varepsilon}{kT}\right) \sigma^3 - \pi m I_2 kT \left(\frac{\partial \rho}{\partial P}\right)_{hc} m^2 \left(\frac{\varepsilon}{kT}\right)^2 \sigma^3 \quad (5)$$

ρ is the density number of the fluids, I_1 and I_2 are perturbation integral. a^{assoc} is calculated by:

$$a^{assoc} = \sum_A \left[\ln X^A - \frac{X^A}{2} \right] + 0.5S$$

(6)

X^A is the fraction of unbounded monomers of single site A and S is the number of associated sites. In present paper, the PC-SAFT model was applied to correlate the density of binary mixtures. In the case of alcohols, five molecular parameters and for xylene, three molecular parameters were fitted to the experimental vapour pressures and densities in order to optimize them. The

PC-SAFT model contains one adjustable parameter k_{ij} which is usually independent of temperature. Values of these fitting parameters are reported in Table 2 along with average absolute deviation (AAD). Results of Table 2 indicate that the PC-SAFT equations is a powerful expression to correlate the density of binary mixtures with only one binary interaction parameter covering the whole concentration range.




Figure 1. Excess molar volumes V_m^E vs. mole fraction of xylene for binary mixtures of xylene with (○) 1-butanol, (◊) 1-pentanol, (●) 1-hexanol, (■) 1-heptanol, (▲) 1-octanol at $T = 323.15$ K.

References

1. Parveen S, Shukla D, Singh S, Singh KP, Gupta M, Shukla JP (2009) *Applied Acoustics*, 70: 507–513
2. Iloukhani H, Almasi M (2011) *Model. J. Solution Chem*, 40: 284–298
3. Almasi M (2016) *J. Therm. Anal. Calorim*, 124: 399–405
4. Almasi M, Khosravi L (2012) *J. Serb. Chem. Soc*, 77: 363–370
5. Redlich OJ, Kister AT (1948) *Ind. Eng. Chem*, 40: 345–348
6. Gross J, Sadowski G (2001) *Ind. Eng. Chem. Res*, 40: 1244–1260
7. Gross J, Sadowski G (2002) *Ind. Eng. Chem. Res.* 41:5510–5515.

How to cite this manuscript: Mohammad Almasi. Theoretical and Experimental Methods for Studying Volumetric Behavior of Binary mixtures Containing Normal Alkanols. Asian Journal of Nanoscience and Materials, 2018, 1, 166-171.